

*cillis capitis lateralibus et linea super nares latè per genas excurrente, colloque anteriori nigris; summo capite et nuchâ vinaceo-cinereis; corpore supra cinereo; remigibus alarum cinereo-fuscis, remige primo externè albo; rectricibus caudæ duabus intermediis fuscis, ad marginem pallidioribus; reliquis nigrescentibus externâ utrinque albo marginatâ; rostro pedibusque nigris.*

Long. tot. unc. 8; alæ,  $4\frac{1}{2}$ ; caud., 3; rost.,  $\frac{4}{5}$ ; tars., 1.

Hab. Erzeroum.

FRINGILLA SANGUINEA. *Fring. brunnea, summo capite nigro; remigum pogoniis externis sanguineo lavatis; primariis nigris; secondariis nigris, ad apices albis; tectricibus caudæ et regione circum-oculari sanguineo lavatis; rectricibus caudæ duabus intermediis nigris, reliquis plus minusve albo notatis, externâ utrinque ferè albâ; rostro flavo; pedibus fuscis.*

Long. tot. 6 unc.  $\frac{3}{4}$ ; alæ, 4; caud.,  $2\frac{1}{2}$ ; rost.,  $\frac{2}{3}$ ; tars.,  $\frac{3}{4}$ .

Hab. Erzeroum.

Mr. Gould afterwards described a new *Ibis* from Hayti, presented by John Hearne, Esq., Corr. Mem., as

IBIS ERYTHRORHYNCHA. *Ib. dorso, alis caudâque metallicè viridibus; capite colloque superiore nigrescenti-cinereis, albo sparsis; corpore sublùs nigrescenti-cinereo; rostro pedibusque rubris; rostro ad basin nigro.*

Hab. Hayti.

Mr. Yarrell exhibited a quill from the wing of a Harpy Eagle, which had died while in the Menagerie of the Earl of Derby, and which was found upon examination to be infested with a great number of a species of *Pediculus*. It appeared that these minute creatures had chosen for their place of retreat the hollow of the large quill-feathers; and the specimens forwarded to Mr. Yarrell by the President were filled with their *exuviae*; two circular apertures situated near the base of the quill afforded the animals access to its interior.

A specimen of the White-bait presented to the Society by Mr. Williams, was exhibited by Mr. Yarrell in order to show the large size sometimes attained by this species; its dimensions were as follow:

Entire length 6 inches.

Depth, measured about midway between the dorsal fin and the extremity of the head, 1 inch 2 lines.

A collection of insects were upon the table which had been collected at Manilla by Mr. Cuming, at whose request Mr. Westwood had furnished the following descriptions of some of the more interesting species for insertion in the Society's Proceedings.

COLLYRIS (COLLIURIS Latr.) FEMORATA (albitarsis Erichs.?)  
*affinis C. Robynsii et lugubris, V. L. C. capite thoraceque violaceis, labro 7-dentato, antennis capite longioribus, fere filiformibus, articulis 3 et 4 annulo apicali, 5<sup>to</sup> in medio, et basi 6<sup>to</sup> et 7<sup>mo</sup> fulvis, palporum labialium articulo basali albo; thorace strigis nonnullis transversis in parte anticâ cum punctis paucis posticis;*

*elytris œneo-cyaneis, valde punctatis, punctis distinctis, versus apicem punctis multo minoribus, apice ipso truncato emarginato; pedibus cyaneis, femoribus ferrugineis, tarsis piceo-nigris, (duobus posticis albidis in mare exceptis.)*

Long. corp. ♂. lin. 6; ♀ lin. 7.

*Obs.* The possession of both sexes of this insect agreeing precisely together, except in size and the colour of the posterior *tarsi*, induces me to give it as a distinct species, since on comparison, its characters will not agree precisely with those of the species previously described by Dejean, MacLeay, Van der Linden, &c.

**THERATES CORACINA.** Erichs. Act. Acad. Cæs. Nat. Cur. 16.

Suppl. *T. nigra nitidissima, elytris nitore sub purpureo nitentibus, his tuberculo basali alteroque minori ante medium; labro, antennarum articulo basali, abdomine, pedibusque luteis, tarso-rum articulo 1<sup>mo</sup> et 2<sup>do</sup> ad apicem, cœteris fuscis, mandibulis lateis, dentibus nigris.*

Long. corp. lin. 9½.

**MORIO ORIENTALIS.** Dejean Species Gen. i. p. 433.

**CLIVINA CASTANEA.** *C. capite thoraceque castaneis, elytris pedibusque pallidioribus. Cl. Fossori dimidio major, capite minori convexo lateribus valde impresso, fronte punto minuto nec stria longitudinali impressa, thorace subquadrato (postice parum latiori) ad angulos posticos impresso, elytris elongatis parallelis sub-punctato-striatis punctisque tribus majoribus in striam tertiam, femoribus anticis crassis, haud dentatis, tibiis anticis extus dentibus tribus elongatis.*

Long. corp. lin. 3½.

**NIGIDIUS LÆVICOLLIS.** *N. niger nitidus, capite supra depresso punctato, thoracis dorso lœvi, lateribus punctatis margineque antico simplici tuberculo minutissimo vix apparenti, elytris inter strias elevatas triplici punctorum impressorum ordine instructis, tibiis anticis 6 dentatis.*

Long. corp. lin. 9.

**PRIONOCERUS CÆRULEIPENNIS.** Perty, Obs. Coleopt. Ind. Orient. p. 33.

**RHYNCHITES MANILLENSIS.** *R. oblongus, cyaneus, nitidus, rostro longiori nigro, punctato, lineis duabus supra ad basin impressis, thorace conico, punctis minutissimis, longitudinaliter haud canaliculato, impressione transversa cum margine postico parallela, elytris magis cœruleis basi-virescentibus punctato-striatis, corpore subtus et femoribus cyaneis, tibiis tarsis et antennis nigris.*

Long. corp. (cum rostro) lin. 3.

**LAMIA PULCELLATOR.** Affinis L. Rubo et Sehestedii. *L. luteo-fusca, sericea, thorace utrinque spina armato punctisque duobus nigris ante medium disci, elytris basi scabris, singulo maculis 4<sup>ta</sup> irregularibus lacteis, scil. 1<sup>ma</sup> majori obliqua marginali, ante medium locata, 2<sup>da</sup> media, ovata prope suturam, 3<sup>ta</sup> obliqua marginali, 4<sup>ta</sup> versus apicem suturæ, maculisque nonnullis minoribus versus apicem elytrorum, in speciminibus variis vari-*

*antibus, et irregularibus, ejusdem coloris; antennis concoloribus,  
apicibus articulorum obscurioribus.*

Long. corp. lin. 11—16.

**EURYCEPHALUS NIGRIPES.** Dej. Catal.

*Cerambyx nigripes.* Oliv. G. 67. Sp. 68. t. 20. f. 149. ♀.

*—maxillosus.* Oliv. G. 67. Sp. 69. t. 20. f. 147. ♂.

Obs. *Species valde varians, maribus, giganteis. In individuis Manillæ macula thoracis discoidalis multum magnitudine variat. In maribus macula communis nigra versus apicem elytrorum rotundata, nec ad apicem nec ad marginem elytrorum extensa. In fœminis vero apicem totum elytrorum occupat.*

Genus COLASPIIS, SCELODONTA Subgenus novum.

*Corpus breve, rugosum.*

*Caput oculis prominentibus fere thoracis latitudine.*

*Labrum parvum quadratum antice marginatum.*

*Mandibulæ parvæ acutæ subtus dente parvo ante apicem armatæ.*

*Maxillæ lobo interno truncato, externo distincto, tenui, longo acuto.*

*Palpi maxillares et labiales apice acutæ.*

*Antennæ longitudine thoracis articulo 2<sup>do</sup> 3<sup>to</sup> breviori, articulis 5 ultimis majoribus.*

*Thorax fere rotundatus, postice parum latior. Sternæ simplicia lata.*

*Elytra brevia obtusa, thorace dimidio latiora.*

*Femora omnia subtus dente armata. Tibiæ 4 posticæ ad apicem externe emarginatæ.*

**SCELODONTA CURCULIONOIDES.** *S. aureo-viridis cyaneo varia, capite viridi ruguloso-punctato vertice lineis tribus impressis antice conjunctis, lateralibus obliquis, antennis nigris, articulis basalibus viridibus; thorace aureo-viridi, utrinque macula oblonga extus lobata, cyanea; transverse anguloso; elytris rugulosis, punctis majoribus in lineas longitudinales irregulariter dispositis, aureo-viridibus fasciis tribus irregularibus suturaque cyaneis.*

Long. corp. lin. 3.

**FORFICULA TARSATA.** *F. nigra, thorace postice et elytris punctis elevatis nitidis; capite nigro sericeo, antennis longis 25-articulatis, pedibus piceis, apice tibiarum tarsisque albidis, forcipe ♂ longissima, fere corporis longitudine gracili, e basi usque ad medium curvata, denticulis nonnullis internis armata; e medio fere ad apicem recta, apice ipso incurvo acuto; segmentis abdominis in ♀ tuberculis minutis elevatis scabris, serieque in singulo ad marginem posticum tuberculorum majorum, denteque obtuso porrecto in singulo segmento utrinque ad angulum lateralem posticum.*

Long. corp. ♂. lin. 7. forcipis, lin. 5. = 12.

**HYMENOTES.** Genus novum e familia Locustidarum Tetrici affine.

*Corpus valde compressum.*

*Caput mediocre, obliquum.*

*Antennæ breves gracillimæ, filiformes, articulo 1<sup>mo</sup> crasso rotundato, 2<sup>do</sup> multo minori, reliquis longitudine sensim crescentibus.*

*Prothorax maximus foliaceus, valde compressus folium aridum exacte referens, supra et ante caput angulariter porrectus, valde elevatus et postice supra abdomen protensus; parte posticâ subtus, pro receptione alarum et abdominis canaliculata, prosternum in collare pro receptione oris formatum.*

*Pedes inter se basi longe distantes, femoribus præsertim posticis foliaceis, tarsis 3-articulatis, articulo 2<sup>ndo</sup> minutissimo. Pulvilli nulli.*

Species 1. *Hymenotes rhombea.* Membracis r. Fabricius, Ent. Syst. 4. 8. 2. p. Syst. Rh. 7. Cicada r. Linn. Syst. Nat. 2. 704. Alatus.

*Habitat in Jamaica. In Mus. Soc. Linn. Lond. olim Banks.*

Species 2. HYMENOTES 3-ANGULARIS. *H. fusca, prothorace subtriangulari, margine, e fronte ad medium integro et curvato, dein ad apicem obliquo, serrato, femoribus anticis vix foliaceis, posticis latioribus, supra irregulariter incisis.*

Long. corp. lin. 5½. Long. prothoracis lin. 8.

FULGORA APICALIS. Westw. Monogr. Fulg. Trans. Soc. Linn. inedit.

MACHAEROTA ENSIFERA. Burmeister, Handb. der Ent. 2. p. 128.

CENTROTUS BIFOLIATUS. *C. fuscus punctatus prothorace antice in cornu longissimum erectum conicum, postice setigerum elevato; apice bifido, parte bifida fere longitudine cornū, retro extensa, cum apice extremo singuli dilatato, acuto; parte postica prothoracis compressa, longitudine corporis apice sensim attenuato et paullo deflexo-curvato, prothorace in medio, supra dorsum, fascia lata albida; hemelytris fuscis punctis nonnullis obliquis pallidis ad marginem internum, tibiis anticis latioribus, posticis gracilibus.*

Long. corp. lin. 2½. Long. e basi usque ad apicem cornu prothoracis, lin. 6.

*Obs.* Valde affinis Centr. Hardwickei, Kirby in Loudon's Mag.

Nat. ii. p. 21. a. e Nepalia.

CENTROTUS HORRIFICUS. *C. fuscus rude punctatus, ab domine nigro, punctis majoribus; prothoracis parte antica in cornua duo elongata erecta divergentia setigera, apice singuli dilatato et acute emarginato, producta; parte postica, abdominis longitudine, basi (supra scutellum distinctum bifidum) in nodum setigerum curvata, nodoque paullo ante apice multo majori elevato-compresso supra rotundato, setigero, armata; tibiis 4 anticis oblongo-ovatis, hemelytrorum venis basalibus tuberculatis.*

Long. corp. lin. 2.

REDUVIUS TIBIALIS. *R. amœno, Guér. (Icon. R. An. Ins. pl. 56. f. 17.) valde affinis, capite postice magis attenuato. Niger, punctatus, thoracis lobo antico parvo tuberculis duobus elevatis armato, hemelytris fuscis, plaga obliqua media, maculaque minori ovata subapicali albido-hyalinis, abdominis lateribus valde dilatatis, antennis, rostro, pedibusque rubris, femoribus nigris apice rubris.*

Long. corp. lin. 12. Expans. alar. lin. 19.