

Molecular differentiation of four *Reptalus* species (Hemiptera: Cixiidae)

S. Bertin^{1†}, L. Picciau^{1†}, Z. Ács², A. Alma¹ and D. Bosco^{1*}

¹DIVAPRA – Entomologia e Zoologia applicate all'Ambiente 'Carlo Vidano', Università di Torino, Via L. da Vinci 44, 10095 Grugliasco (TO), Italy; ²Fitolab Plant Pest Diagnostic and Advisory Ltd. Istenhegyi út 29, Budapest H-1125, Hungary

Abstract

The cixiid species *Reptalus quinquecostatus*, *R. cuspidatus*, *R. panzeri* and *R. melanochaetus* are widely distributed in Europe and are receiving growing attention because of their potential role as phytoplasma vectors. Identifying the *Reptalus* species is restricted to a few specialist entomologists and relies on the morphology of the male genitalia, hampering the identification of juveniles and females. This study provides the tools for species discrimination by integrating the morphological description, which is primarily for the genus identification, with new molecular assays, based on both ribosomal and mitochondrial DNA. PCR-RFLP assays carried out on the mitochondrial cytochrome oxidase I gene (COI) with *AluI* provided species-specific profiles for the four *Reptalus* species. Amplification of a ribosomal internal transcribed spacer (ITS2) region produced species-specific fragments of different sizes for *R. quinquecostatus*, *R. melanochaetus*, *R. cuspidatus* and *R. panzeri*. The digestion of the ITS2 PCR product with *TaqI* allowed the discrimination of these latter two species. This molecular identification key ensures reliable results and can be successfully applied not only to adults, but also to the nymphs feeding on the roots. The identification of the nymphs (i) extends the collection period of these monovoltine species to the whole year (adults are present for a short summer period) and (ii) allows the unambiguous identification of their actual host plants because nymphs are steady on the root system while adults tend to disperse onto other plants. Fast and reliable identification of the *Reptalus* species provides useful help in monitoring activities and, therefore, in designing rational control strategies to protect crops from phytoplasma infection.

Keywords: Cixiidae, species identification, ITS2, mtCOI, PCR-RFLP, DNA barcoding

(Accepted 8 October 2009)

Introduction

The genus *Reptalus* (Hemiptera: Cixiidae) includes 11 recognised species distributed in the temperate areas of central

and southern Europe and the Mediterranean basin (Hoch, 2004 available online at 'http://www.faunaeur.org').

Though phylogenetic relationships between *Reptalus* and the other cixiid genera have been recently clarified (Ceotto & Bourgoïn, 2008; Ceotto *et al.*, 2008), the biology and behaviour of the *Reptalus* species remain poorly known because of the peculiar characteristics of this planthopper family. The lifecycle of cixiids is closely related to their host plants, which are exploited for feeding, mating and oviposition and serve as protection against predators (Wilson *et al.*,

*Author for correspondence

Fax: 39 011 6708535

E-mail: domenico.bosco@unito.it

† The first two authors contributed equally to the paper.

Table 1. Field-collected samples and their host plants.

Species (accession numbers)	Sampling location	Sample size	Host plants
<i>R. quinquecostatus</i> (ITS2: GQ390297) (COI: GQ397852)	Piemonte (Italy)	26	<i>Ulmus, Malus, Salix, Rubus, Sinapis, Convolvulus</i>
	Emilia Romagna (Italy)	10	Vineyard weeds
	Azov (South Russia)	2	Potato field
	Csíksereda (Romania)	3	Potato field
	Fundulea (Romania)	13	Potato field
<i>R. cuspidatus</i> (ITS2: GQ390298) (COI: GQ397853)	Andornaktalya (Hungary)	4	Vineyard weeds
	Piemonte (Italy)	57	<i>Artemisia, Erodium, Viola, Convolvulus, Calystegia, Rubus, Sinapis</i>
	Valle d'Aosta (Italy)	19	<i>Convolvulus</i>
	Emilia Romagna (Italy)	1	Vineyard weeds
	<i>R. panzeri</i> (COI: GQ397854) (ITS2: GQ390299)	Piemonte (Italy)	4
Toscana (Italy)		1	<i>Ulmus</i>
Veneto (Italy)		1	<i>Ulmus</i>
Emilia Romagna (Italy)		1	<i>Clematis</i>
Fundulea (Romania)		9	Potato field
Andornaktalya (Hungary)		6	Vineyard weeds
<i>R. melanochaetus</i> (ITS2: GQ390300) (COI: GQ397855)	Piemonte (Italy)	4	<i>Prunus, Ulmus</i>
	Emilia Romagna (Italy)	2	<i>Convolvulus</i>

1994; Sforza & Bourgoïn, 1998). Adult females lay eggs in the soil near the base of a host plant, and the five nymphal instars live underground and feed on its roots. Newly emerged adults fly to leaves and shoots for feeding and mating (Wilson & Tsai, 1982; O'Brien & Wilson, 1985). A broad host-plant range has been identified, and the available data indicate that different cixiid species belonging to the same genus may feed on distantly related plant families (Wilson *et al.*, 1994). The cixiids are mostly phloem-feeders, and their sap-sucking habit makes them potential vectors of phloem-inhabiting plant pathogens. Indeed, several species are known as vectors of viruses, bacterium-like organisms and phytoplasmas (Julia *et al.*, 1985; Danet *et al.*, 2003; Weintraub & Beanland, 2006). In Europe, several crops (beet, maize, strawberry, solanaceous and grapevine) are affected by different diseases vectored by cixiids (Sforza *et al.*, 1998; Danet *et al.*, 2003; Březíková & Linhartová, 2007; Jović *et al.*, 2007; Bressan *et al.*, 2008).

Within the genus *Reptalus*, some species have recently been noted as potential vectors of phytoplasmas belonging to the stolbur group (*Candidatus* Phytoplasma solani, 16SrXII-A genetic group). Individuals of *Reptalus panzeri* (Löw) collected in maize fields affected by maize redness, a disease recently associated to a phytoplasma infection (Duduk & Bertaccini, 2006), tested positive for stolbur in PCR assays. In further studies, the efficiency of this species to transmit stolbur to healthy maize plants was confirmed by transmission trials in mesh cages (Jović *et al.*, 2007). Some *Reptalus* species are also suspected to be vectors of a serious grapevine yellow, known as bois noir (BN). BN is an endemic stolbur disease of grape-growing areas in Europe, Asia Minor and the Mediterranean basin and has shown an increasing economic importance over the past years (Maixner, 2006). Although the only confirmed vector of BN is the cixiid *Hyalesthes obsoletus* (Signoret), *R. panzeri* and *Reptalus quinquecostatus* (Dufour) collected in affected vineyards have been found positive to stolbur phytoplasma (Palermo *et al.*, 2004; Trivellone *et al.*, 2005; Riedle-Bauer

et al., 2006). It has also been demonstrated that *R. quinquecostatus* can successfully inoculate BN into an artificial feeding medium through parafilm (Pinzauti *et al.*, 2008). The potential involvement of *Reptalus* in phytoplasma transmission has prompted an active monitoring practice. The epidemiological studies and application of control strategies rely on careful surveys of the spread of cixiid species. The reliable identification of each species becomes essential, but may sometimes produce uncertain results. Identification of cixiids, including *Reptalus*, is based on morphological characteristics and restricted to a small number of specialist entomologists with broad experience of this insect family. Even for experts, the morphological distinction of closely related species remains difficult. Furthermore, the main taxonomic characteristics concern the male genitalia, thus hampering the identification of juveniles and adult females.

DNA-based approaches offer valuable support to traditional taxonomic methods and nowadays are widely employed for insect species identification. Considering both chromosomal and extra-chromosomal DNA, the ribosomal and mitochondrial regions have proven informative for taxonomic purposes. Notably, the internal transcribed spacer regions (ITS1 and 2) of the ribosomal operons are highly polymorphic and support phylogenetic analyses at species level (Li, 2007; Roe & Sperling, 2007). Among the mitochondrial sequences, the cytochrome oxidase I gene (COI) is frequently employed (Behere *et al.*, 2008; Sinclair & Gresens, 2008) and has been chosen as the reference point for DNA barcoding, a universal system of animal species identification. DNA barcoding is based on the premise that sequence diversity within a short, standardised segment of the genome (on average 600–800 bp) can provide a 'biological barcode' that enables identification to the species level (Hebert *et al.*, 2003; Hebert & Gregory, 2005). Previous studies have shown that the COI polymorphism can effectively resolve species recognition (Hebert *et al.*, 2004; Hogg & Hebert, 2004; Smith *et al.*, 2005).




Fig. 1. External morphological features of *Reptalus* genus: (a) head and thorax; t, tegula; p, pronotum; m, mesothorax with five evident carinae; (b) tip of a forewing showing the veins with tubercles; (c) head of *R. quinquecostatus* in frontal view; a, antenna; ac, anteclypeus; e, eye; f, front; oc, ocelli; pc, postclypeus; v, vertex; (d) detail of vertex; tk, transverse keels; (e) hind tarsus of *R. cuspidatus* (subgenus *Trepalus*) showing the first segment with spines bearing platellae.

The aim of this study was to provide a molecular identification key of the most common European *Reptalus* species (Holzinger *et al.*, 2003; Hoch, 2004). We tested the reliability of both ITS2 and COI DNA polymorphisms for the identification of four species, *R. quinquecostatus* (subgenus *Reptalus*), *Reptalus cuspidatus* (Fieber) (subgenus *Trepalus*), *R. panzeri* (subgenus *Reptalus*) and *Reptalus melanochaetus* (Fieber) (subgenus *Reptalus*) and correlated morphological characteristics of the different species with species-specific DNA profiles.

Material and methods

Samples collection

Adults of *R. quinquecostatus* ($n=58$), *R. cuspidatus* ($n=77$), *R. panzeri* ($n=22$) and *R. melanochaetus* ($n=6$) were field-collected from different Italian and eastern European regions during the summers of 2007 and 2008. Field surveys were mainly targeted at those plants known as potential hosts of the *Reptalus* species (Holzinger *et al.*, 2003; Mazzoni, 2005) and those potentially infected by stolbur phytoplasma (table 1).

Specimens of both sexes were preserved in absolute ethanol until their morphological identification and/or DNA extraction.

Morphological species identification

Specimens were individually identified with a stereo microscope and the external morphological features were

used to determine family and genus. For species identification, male genitalia (aedeagus, parameres and anal tube) were carefully dissected and placed in a 10% potassium




Fig. 2. *Reptalus quinquecostatus*: habitus in dorsal view. Scale bar: 1 mm.




Fig. 3. Aedeagi of the four *Reptalus* species: (a) *R. quinquecostatus*, ventral view. (b) *R. cuspidatus*, ventral view. (c) *R. panzeri*, ventral view. (d) *R. melanochaetus*, ventral view. Scale bar: 0.5 mm.

hydroxide solution for about one day in order to remove membranous soft tissues. This procedure also turns the male genitalia semi-diaphanous, which allows better visualisation of all the shape details (H. Hoch, personal communication).

DNA extraction

Total genomic DNA was extracted from individual planthoppers following the protocol described in Bosco *et al.* (2002). Briefly, the ethanol-preserved adults were dried onto filter paper and homogenised in a CTAB-based buffer (2% w/v cetyl-trimethyl-ammonium-bromide (CTAB); 1.4 M NaCl; 20 mM EDTA pH 8.0; 100 mM Tris-HCl pH 8.0; 0.2% mercaptoethanol). The male specimens were homogenised after removal of the aedeagus for morphological observation and species identification. After incubation at 65°C for 30 min, DNA was extracted with one volume of chloroform:isoamylalcohol 24:1 solution and then precipitated with the addition of two volumes of absolute ethanol. The DNA pellet was then washed with 70% ethanol, vacuum dried and resuspended in 100 µl TE pH 8.0.

PCR and sequencing analyses

The amplification of the ITS2 region was carried out with the primers ITS2fw (5'-TGTGAACTGCAGGACACATG-3') and ITS2rv (5'-ATGCTTAAATTTAGGGGGTA-3'), which respectively anneal on 5.8S and 28S ribosomal regions (Collins & Paskewitz, 1996). PCR was performed in 25 µl reaction volume containing: 1 × PCR buffer (20 mM Tris-HCl pH 8.4, 50 mM KCl), 1.5 mM MgCl₂, 200 µM of each dNTP, 0.5 µM of each primer, 1 unit of *Taq* polymerase (Invitrogen, Carlsbad, CA) and 50–150 ng of template DNA. The thermocycling conditions consisted of an initial denaturation cycle at 94°C for 5 min, 35 cycles at 94°C for 30 s, 58°C for 45 s and 72°C for 1 min and a final cycle at 72°C for 10 min.

A fragment of COI mitochondrial gene was amplified using the primers C1-J-2195 (5'-TTGATTTTTTGGTCATC-CAGAAGT-3') and TL2-N-3014 (5'-TCCAATGCACTAAT-CTGCCATATTA-3') (Simon *et al.*, 1994). The PCR conditions were the same as described for ITS2 amplification, except the MgCl₂ content (increased to 2.5 mM) and the annealing step (54°C for 1 min).

Products of both PCR assays were separated by electrophoresis onto a 1% (w/v) agarose gel and then directly purified from the amplification tube using the PureLink™




Fig. 4. (a) agarose gel electrophoresis of ITS2 PCR products from the four *Reptalus* species. *R.q.*, *R. quinquecostatus*; *R.c.*, *R. cuspidatus*; *R.p.*, *R. panzeri*; *R.m.*, *R. melanochaetus*; M: molecular weight marker (1 kb ladder); (b) RFLP profile after *TaqI* digestion of the ITS2-PCR product for the two species *R. cuspidatus* and *R. panzeri*. *R.c.*: *R. cuspidatus*; *R.p.*: *R. panzeri*; M: molecular weight marker (100 bp ladder).

PCR Purification Kit (Invitrogen, Carlsbad, CA, USA). The ITS2 and COI purified products were sequenced (Bio-Fab Research, Pomezia, Italy) in both directions using the respective forward and reverse primers. The mitochondrial and ribosomal consensus sequences of each *Reptalus* species have been deposited in GenBank (National Centre for Biotechnology Information, NCBI). Accession numbers for ITS2 sequences: GQ390297–GQ390300; accession numbers for COI sequences: GQ397852–GQ397855 (table 1). COI sequences are also available in Barcode of Life databank.

RFLP analysis

Restriction sites on both ITS2 and COI sequences were predicted using Webcutter 2.0 (available online at 'http://ma.lundberg.gu.se/cutter2/'). The ITS2 fragments amplified

from the *R. cuspidatus* and *R. panzeri* species were digested with the *TaqI* restriction enzyme at 65°C for 2 h. COI amplicons from all tested species were subjected to *AluI* restriction activity at 37°C for 2 h.

For both ITS2 and COI samples, 5 µl of PCR product was digested with one unit of the proper enzyme in a 15 µl reaction volume, according to the manufacturer's instructions (Invitrogen, Carlsbad, CA, USA). The restriction fragments were separated by electrophoresis on 2% (w/v) 1 × TBE agarose gels (MetaPhore agarose for resolution of small nucleic acids; Cambrex, Rockland, ME, USA) at 70 V for 2 h and stained with ethidium bromide.

Results

Morphological species identification

A brief review of the external characteristics for identifying cixiids among the different Auchenorrhyncha taxa is given. The main features of the infraorder Fulgoromorpha are (i) the presence of a tegula at the base of each forewing (fig. 1a) and (ii) two typically 'Y-shaped' anal veins on the clavus. A short anteclypeus and the median coxae clearly separated at the base are also present.

The family Cixiidae is quite easily distinguishable from the other Fulgoromorpha by the large and transparent forewings, surpassing the tip of the abdomen, with distinct venations, often with tubercles (fig. 1b). These forewings are held roof-like in the rest position and sometimes overlap slightly apically. The frons is wide, with three keels giving a distinctive aspect to the species belonging to this family. Three ocelli are normally present, two laterally and one in the middle on the frons (fig. 1c). The pronotum is short and collar-like. The mesonotum is quite large, subtriangular and bears three or five longitudinal carinae (fig. 1a). The hind tibiae have a row of apical spines and no movable spurs. The genus *Reptalus* differs from the other genera principally by bearing five distinct longitudinal carinae on the mesonotum (fig. 1a), and two transverse keels on the head (fig. 1d). The size of the body normally exceeds 5 mm in length. The first segment of the hind tarsus bears apical spines with platellae (macrochaetae) in the subgenus *Trepalus* (fig. 1e); platellae are absent in the subgenus *Reptalus*. Finally, the head in lateral view appears obtuse (Ossiannilsson, 1978; Holzinger *et al.*, 2003; Wilson, 2005). An overview of the insect body is given for *R. quinquecostatus* (fig. 2).

The *Reptalus* species identification mainly relies on the shape of the aedeagus. The drawings representing the aedeagi of the four considered species are provided in fig. 3 in order to show the distinctive features. Parameters and anal tube are omitted.

ITS2 sequence: PCR and RFLP analyses

The ITS2 region was successfully amplified from each of the four *Reptalus* species, providing fragments with the following sizes: 1450 bp for *R. quinquecostatus*; 850 bp for *R. cuspidatus* and *R. panzeri*; and 960 bp for *R. melanochaetus* (fig. 4a). The four species shared a similar sequence within the first 200 bp and the last 550 bp of the ITS2 region (range of similarity scores: 89–93%); a diverse number of repeats occurred between these two ends, leading to different fragment lengths.




Fig. 5. *AluI* restriction sites of the four *Reptalus* species.

The sequence analysis revealed no polymorphism among the individuals belonging to the same species and, consequently, no differences amenable to the geographical origin, host plant or sex were noticed. The absence of polymorphism at the intra-specific level led us to exclude any variability among the different ribosomal units within the same individual genome.

The ITS2 amplicon length polymorphism did not enable us to distinguish between the *R. cuspidatus* and *R. panzeri* species. In this case, the correct species identification was achieved by further performing a RFLP assay. The *TaqI* digestion produced four restriction fragments for *R. panzeri* (~20 bp, ~230 bp, ~260 bp, ~340 bp), while no *TaqI* restriction sites were present within the *R. cuspidatus* sequence (fig. 4b).

COI sequence: PCR and RFLP analyses

Amplification of the mitochondrial COI gene from all tested individuals always provided a 920 bp fragment. The alignment of the COI sequences revealed a certain degree of polymorphism among the four different species, while no variability was recorded within the same species. Polymorphisms were found at the *AluI* recognition site (AGCT), generating species-specific digestion patterns (fig. 5). According to the mutation points revealed by sequencing, the endonuclease activity produced three restriction fragments for *R. quinquecostatus* (~170 bp, ~260 bp, ~490 bp) and *R. cuspidatus* (~80 bp, ~330 bp, ~510 bp), two restriction fragments for *R. panzeri* (~440 bp, ~480 bp) and four restriction fragments for *R. melanochaetus* (~60 bp, ~120 bp,

~170 bp, ~570 bp) (fig. 6). The smallest fragments (60 bp for *R. melanochaetus* and 80 bp for *R. cuspidatus*), even though faint, could be detected by agarose gel electrophoresis. The RFLP patterns were constant within each species, confirming the absence of variability at an intra-specific level.




Fig. 6. RFLP profile after *AluI* digestion of the COI-PCR product for the four *Reptalus* species. *R.q.*, *R. quinquecostatus*; *R.c.*, *R. cuspidatus*; *R.p.*, *R. panzeri*; *R.m.*, *R. melanochaetus*; M: molecular weight marker (100 bp ladder).

Discussion

This study provides a comprehensive description of both morphological and molecular tools for species discrimination within the genus *Reptalus*. *R. cuspidatus* and *R. panzeri* were abundant in Italy and eastern Europe, respectively, while *R. quinquecostatus* was widespread in all the considered areas. *R. melanochaetus* was shown to be less represented in the investigated areas, according to Mazzoni (2005), who did not report this species in his recent surveys of Auchenorrhyncha in Tuscany, Central Italy.

Two different DNA regions, ribosomal ITS2 and mitochondrial COI, allowed discrimination at the species level. Both sequences diverged among the four *Reptalus* species and were suitable for correctly assigning the individuals to each species in accordance with the previous morphological identification. The absence of within-species polymorphisms proves the species-specificity of both ITS2 and COI markers.

The ITS2 PCR allowed us to recognise two of the four *Reptalus* species directly by amplicon size polymorphism. A further RFLP analysis was required when an 850bp fragment, corresponding to both the *R. cuspidatus* and *R. panzeri* species, was amplified. Despite their length polymorphism, ITS2 sequences possess similar regions in all of the four species. It is possible that such conserved domains were part of a precursor rRNA and have been differently truncated or extended during rDNA duplication. This could have produced diverse sequences or alleles then become fixed separately during speciation (Koekemoer *et al.*, 2009).

Hence, the ITS2 region provided the simplest tool for rapid *Reptalus* identification because the PCR assay was sufficient for recognising at least two species. Unambiguous species discrimination also can be achieved by routinely performing RFLP assays on COI-PCR products. In this case, the sole *AluI* enzyme allowed clear differentiation of the four species with a unique restriction reaction.

The reliability of the ITS2 and COI markers has been tested on different populations from different sites in Italy and eastern Europe and no polymorphisms ascribable to the geographic origins have been noticed. The two regions also showed stable PCR or RFLP patterns irrespective of the different plants hosting the insects. Given their conservation and specificity, both ribosomal and mitochondrial regions can be considered potential identification keys. These new tools require rapid and handy procedures and may contribute to broadening the current handful of competent scientists in cixiid identification. Furthermore, the variation in *Reptalus* COI gene should be considered for future DNA barcode analyses on cixiid species. The correspondence between morphological and COI-based identifications suggests that this short DNA fragment can be a good identifier for this insect family. The presence of cixiids in barcoding databases has been restricted to very few species, and the uploading of *Reptalus* COI sequences may enrich this universal and easily accessible identification system.

Both COI and ITS2 sequences can be efficiently applied to the species identification of juveniles and females, for which a morphological identification tool is unavailable. From this perspective, this work represents a useful contribution to faunal studies and insect vector monitoring. First, the ability to correctly identify nymphs allows the collection period to be extended to the whole year, making possible a longer monitoring activity from the current couple of months when

adults are present. Second, while adults tend to rapidly disperse and can have a more polyphagous feeding habit, nymphs are harboured exclusively on the root system, making feasible the unambiguous association with their actual host plants. So far, breeding plants of *R. cuspidatus*, *R. melanochaetus*, *R. panzeri* and *R. quinquecostatus* are unknown and, during our surveys, we were able to definitely associate *R. cuspidatus* with *Artemisia verlotiorum* Lamotte since many larvae were found on its roots.

The correct identification of the *Reptalus* host plants, together with the identification of plants that can act as a reservoir of plant pathogens for crop species, is of fundamental importance in designing a rational control strategy. The same total DNA preparations from *Reptalus* spp. can serve as a PCR template for insect species and stolbur phytoplasma identification (data not shown).

In conclusion, this study provides fast and reliable assays for the correct identification of four *Reptalus* species. Morphological features are suggested for genus identification while species discrimination within the genus can be easily achieved with ITS2 and/or COI sequence analysis. The integration of morphological and molecular approaches provides useful help in monitoring activities and the rational control of potential vector species.

Acknowledgements

The authors would like to thank all the people who kindly helped in sample collection, particularly Mirko Bacchiavini and Pasquale Mazzio (Consorzio Fitosanitario Provinciale di Reggio Emilia), Ruggero Mazzilli, Paolo Fontana and Mauro Daccordi. The authors also thank Federica Tota for helping in laboratory analysis and Prof. Hannelore Hoch for her precious advices. This work was supported by the Piemonte Region and the MIUR-PRIN 2008.

References

- Behere, G.T., Tay, W.T., Russell, D.A. & Batterham, P. (2008) Molecular markers to discriminate among four pest species of *Helicoverpa* (Lepidoptera: Noctuidae). *Bulletin of Entomological Research* **98**, 599–603.
- Bosco, D., Palermo, S., Mason, G., Tedeschi, R., Marzachi, C. & Boccardo, G. (2002) DNA-based methods for the detection and the identification of phytoplasmas in insect vector extracts. *Molecular Biotechnology* **22**, 9–18.
- Bressan, A., Sémétey, O., Nusillard, B., Clair, D. & Boudon-Padieu, E. (2008) Insect vectors (Hemiptera: Cixiidae) and pathogens associated with the disease syndrome 'basses richesses' of sugar beet in France. *Plant disease* **92**, 113–119.
- Březíková, M. & Linhartová, S. (2007) First report of potato stolbur phytoplasma in Hemipterans in southern Moravia. *Plant Protection Science* **43**, 73–76.
- Ceotto, P.C. & Bourgoïn, T. (2008) Insights into the phylogenetic relationships within Cixiidae (Hemiptera: Fulgoromorpha): cladistic analysis of a morphological data set. *Systematic Entomology* **33**, 484–500.
- Ceotto, P.C., Kergoat, G.J., Rasplus, J.Y. & Bourgoïn, T. (2008) Molecular phylogenetics of cixiid planthoppers (Hemiptera: Fulgoromorpha): New insights from combined analyses of mitochondrial and nuclear genes. *Molecular Phylogenetics and Evolution* **48**, 667–678.

- Collins, F.H. & Paskewitz S.M. (1996) A review of the use of ribosomal DNA (rDNA) to differentiate among cryptic *Anopheles* species. *Insect Molecular Biology* **5**, 1–9.
- Danet, J.L., Foissac, X., Zreik, L., Salar, P., Verdin, E., Nourrisseau, J.G. & Garnier, M. (2003) 'Candidatus *phlomobacter fragariae*' is the prevalent agent of marginal chlorosis of strawberry in French production fields and is transmitted by the planthopper *Cixius wagneri* (China). *Phytopathology* **93**, 644–649.
- Duduk, B. & Bertaccini, A. (2006) Corn with symptoms of reddening: new host of stolbur phytoplasma. *Plant Disease* **90**, 1313–1319.
- Hebert, P.D.N. & Gregory, T.R. (2005) The promise of DNA barcoding for taxonomy. *Systematic Biology* **54**, 852–859.
- Hebert, P.D.N., Cywinka, A., Ball, S.L. & deWaard, J.R. (2003) Biological identification through DNA barcodes. *Proceedings of the Royal Society of London, Series B* **270**, 313–321.
- Hebert, P.D.N., Penton, E.H., Burns, J.M., Janzen, D.H. & Hallwachs, W. (2004) Ten species in one: DNA barcoding reveals cryptic species in neotropical skipper butterfly *Astraptes fulgerator*. *Proceedings of the National Academy of Science* **101**, 14812–14817.
- Hoch, H. (2004) Fauna Europaea: Cixiidae, Reptalus. Fauna Europaea version 1.1, available online at <http://www.faunaeur.org>.
- Hogg, I.D. & Hebert, P.D.N. (2004) Biological identification of springtails (Hexapoda: Collembola) from the Canadian Arctic, using mitochondrial DNA barcodes. *Canadian Journal of Zoology* **82**, 749–754.
- Holzinger, W.E., Kammerlander, I. & Nickel, H. (2003) *The Auchenorrhyncha of Central Europe – Die Ziekaden Mitteleuropas Volume 1: Fulgoromorpha, Cicadomorpha excl. Cicadellidae*. 673 pp. Leiden, The Netherlands, Brill.
- Jović, J., Cvrković, T., Mitrović, M., Krnjajić, S., Redinbaugh, M.G., Pratt, R.C., Gingery, R.E., Hogenhout, S.A. & Toševski, I. (2007) Roles of stolbur phytoplasma and *Reptalus panzeri* (Cixiinae, Auchenorrhyncha) in the epidemiology of Maize redness in Serbia. *European Journal of Plant Pathology* **118**, 85–89.
- Julia, J.F., Dollet, M., Randles, J. & Calvez, C. (1985) Foliar decay of coconut by *Myndus taffini* (FDMT): new results. *Oléagineux* **40**, 19–27.
- Koekemoer, L.L., Misiani, E.A., Hunt, R.H., Kent R.J., Norris, D.E. & Coetzee, M. (2009) Cryptic species within *Anopheles longipalpis* from southern Africa and phylogenetic comparison with members of the *An. funestus* group. *Bulletin of Entomological Research* **99**, 41–49.
- Li, Z.-X. (2007) Molecular differentiation of the four most commonly occurring *Trichogramma* (Hymenoptera: Trichogrammatidae) species in China. *European Journal of Entomology* **104**, 363–367.
- Maixner, M. (2006) Grapevine yellows – Current developments and unsolved questions. In *Proceedings of the 15th meeting of the International Council for the Study of Virus and Virus-like Diseases of the Grapevine (ICVG), Stellenbosch, South Africa, 3–7 April 2006*. Available online at <http://www.icvg.ch/data/maixner.pdf>.
- Mazzoni, V. (2005) Contribution to the knowledge of the Auchenorrhyncha (Hemiptera Fulgoromorpha and Cicadomorpha) of Tuscany (Italy). *Redia* **88**, 85–102.
- O'Brien, L.B. & Wilson, S.W. (1985) Planthopper systematics and external morphology. pp. 61–102 in Nault, L.R. & Rodriguez, J.G. (Eds) *The Leafhoppers and Planthoppers*. New York, NY, USA, John Wiley and Sons.
- Ossiannilsson, F. (1978) *The Auchenorrhyncha (Homoptera) of Fennoscandia and Denmark. Part 1: introduction, infraorder Fulgoromorpha*. 222 pp. Klampenborg, Denmark, Scandinavian Science Press Ltd.
- Palermo, S., Elekes, M., Botti, S., Ember, I., Alma, A., Orosz, A., Bertaccini, A. & Kölber, M. (2004) Presence of stolbur phytoplasma in Cixiidae in Hungarian vineyards. *Vitis* **43**, 201–203.
- Pinzauti, F., Trivellone, V. & Bagnoli, B. (2008) Ability of *Reptalus quinquecostatus* (Hemiptera: Cixiidae) to inoculate stolbur phytoplasma to artificial feeding medium. *Annals of Applied Biology* **153**, 299–305.
- Riedle-Bauer, M., Tiefenbrunner, W., Otreba, J., Hanak, K., Schildberger, B. & Regner, F. (2006) Epidemiological observations on bois noir in Austrian vineyards. *Mitteilungen Klosterneuburg, Rebe und Wein, Obstbau und Früchteerwertung* **56**, 166–170.
- Roe, A.D. & Sperling, F.A.H. (2007) Population structure and species boundary delimitation of cryptic *Dioryctria* moths: an integrative approach. *Molecular Ecology* **16**, 3617–3633.
- Sforza, R. & Bourgoin, T. (1998) Female genitalia and copulation of the planthopper *Hyalesthes obsoletus* Signoret (Hemiptera, Fulgoromorpha, Cixiidae). *Annales de la Société Entomologique de France (N.S.)* **34**, 63–70.
- Sforza, R., Clair, D., Daire, X., Larrue, J. & Boudon-Padieu, E. (1998) The role of *Hyalesthes obsoletus* (Hemiptera: Cixiidae) in the occurrence of bois noir of grapevines in France. *Journal of Phytopathology* **146**, 549–556.
- Simon, C., Frati, F., Beckenbach, A., Crespi, B., Liu, H. & Flook, P. (1994) Evolution, weighting and phylogenetic utility of mitochondrial gene sequences and a compilation of conserved polymerase chain reaction primers. *Annals of Entomological Society of America* **87**, 651–701.
- Sinclair, C.S. & Gresens, S.E. (2008) Discrimination of *Cricotopus* species (Diptera: Chironomidae) by DNA barcoding. *Bulletin of Entomological Research* **98**, 555–563.
- Smith, M.A., Fisher, B.L. & Hebert, P.D.N. (2005) DNA barcoding for effective diversity assessment of a hyperdiverse arthropod group: the ants of Madagascar. *Philosophical Transactions of the Royal Society of London, Series B* **360**, 1825–1834.
- Trivellone, V., Pinzauti, F. & Bagnoli, B. (2005) *Reptalus quinquecostatus* (Dufour) (Auchenorrhyncha Cixiidae) as a possible vector of Stolbur-phytoplasma in a vineyard in Tuscany. *Redia* **88**, 103–108.
- Weintraub, P.J. & Beanland, L. (2006) Insect vectors of phytoplasmas. *Annual Review of Entomology* **51**, 91–111.
- Wilson, S.W. (2005) Keys to the families of Fulgoromorpha with emphasis on planthoppers of potential economic importance in the south-eastern United States (Hemiptera: Auchenorrhyncha). *Florida Entomologist* **88**, 464–481.
- Wilson, S.W. & Tsai, J.H. (1982) Descriptions of the immature stages of *Myndus crudus* (Homoptera: Fulgoroidea: Cixiidae). *Journal of the New York Entomological Society* **90**, 166–175.
- Wilson, S.W., Mitter, C., Denno, R.F. & Wilson, M.R. (1994) Evolutionary patterns of host plant use by delphacid planthoppers and their relatives. pp. 7–45 in Denno, R.F. & Perfect, T.J. (Eds) *Planthoppers: Their Ecology and Management*. New York, NY, USA, Chapman and Hall.